

VYVGART[®]
(efgartigimod alfa-fcab)
Injection for Intravenous Use
400 mg/20 mL vial


VYVGART Infusions

Information on the infusion process:
before, during, and after

My VYVGART[®] *Path*

VYVGART is given in treatment cycles with an individualized break between cycles*

A treatment cycle consists of a 1-hour infusion each week for 4 weeks (4 infusions per cycle)



Breaks between cycles are individualized because your next cycle of treatment is based on an evaluation of your generalized myasthenia gravis (gMG) symptoms and side effects by your neurologist.* Track your symptoms to help your neurologist determine your next treatment cycle. The safety of starting the next treatment cycle sooner than 4 weeks from the last treatment of the previous treatment cycle has not been established.

*If additional cycles are needed.

What is VYVGART® (efgartigimod alfa-fcab)?

VYVGART is a prescription medicine used to treat a condition called generalized myasthenia gravis, which causes muscles to tire and weaken easily throughout the body, in adults who are positive for antibodies directed toward a protein called acetylcholine receptor (anti-AChR antibody positive).

SELECT IMPORTANT SAFETY INFORMATION

VYVGART may increase the risk of infection. In a clinical study, the most common infections were urinary tract and respiratory tract infections.

Please see Important Safety Information on last page. Please see full Prescribing Information.

YOUR STEP-BY-STEP INFUSION GUIDE

When you begin any new treatment, it's good to be prepared. Reviewing this checklist may help you plan for your visits to your infusion location.

☒ **Check the boxes** as you prepare for your infusion

THE DAY BEFORE YOUR INFUSION, WE RECOMMEND THAT YOU:

- ☐ **Drink plenty of water** and eat how you normally would
- ☐ **Try** to get a full night of sleep
- ☐ **Make a list** of your current medications to bring to your appointment
- ☐ **Plan** how you are getting to the infusion location. You may need to find a ride if you cannot drive yourself

SELECT IMPORTANT SAFETY INFORMATION

More patients on VYVGART vs placebo had below normal levels for white blood cell counts, lymphocyte counts, and neutrophil counts. The majority of infections and blood side effects were mild to moderate in severity. Your health care provider should check you for infections before starting treatment, during treatment, and after treatment with VYVGART.

Please see Important Safety Information on last page. Please see full Prescribing Information.

ON THE DAY OF YOUR INFUSION, WE RECOMMEND THAT YOU:

- ☐ **Wear** comfortable clothes and bring layers in case it's chilly
- ☐ **Drink** plenty of water
- ☐ **Eat** how you normally would
- ☐ Get to your appointment at least **15 minutes early**

DURING EACH INFUSION, A HEALTHCARE PROFESSIONAL WILL:



**Prepare
VYVGART**
for infusion



**Check your vital
signs**, including
your pulse, blood
pressure, and
temperature



Find a vein
in your arm and
put in the
intravenous
(IV) line



Monitor you
during and after
your infusion

You may want something to **pass the time** during the infusion process. Consider bringing a:



Book



Crossword



TV show



Podcast



Game

IMMEDIATELY AFTER YOUR INFUSION APPOINTMENT:

- ☐ **Your infusion may take approximately 1 hour.** During this time and for an hour after your infusion, a doctor or nurse will monitor you for any reactions
- ☐ **You will need 4 infusions total to complete the cycle,** so make sure to book your next infusion if you have not done so already

AFTER YOUR INFUSION:

- ☐ **Carefully track your symptoms.** To do this, you may want to regularly complete a Myasthenia Gravis Activities of Daily Living (MG-ADL) assessment. A My VYVGART Path Nurse Case Manager can teach you how to do this
- ☐ **Talk with your doctor** about how you are feeling
- ☐ **Tell your healthcare provider** immediately about any undesirable reactions, such as rashes, swelling under the skin, and shortness of breath

Still have questions about your VYVGART treatment?

Speak with your doctor if you have any questions or if you want more information about VYVGART.

If you are enrolled in My VYVGART Path, reach out to a Nurse Case Manager for support at **1-833-MY-PATH-1** (1-833-697-2841).

Not yet enrolled in My VYVGART Path?

**ASK YOUR
DOCTOR
TO SIGN YOU UP**



With My VYVGART Path, a Nurse Case Manager is waiting to help you through your VYVGART treatment journey.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about VYVGART® (efgartigimod alfa-fcab)?

VYVGART may cause serious side effects, including:

- **Infection.** VYVGART may increase the risk of infection. In a clinical study, the most common infections were urinary tract and respiratory tract infections. More patients on VYVGART vs placebo had below normal levels for white blood cell counts, lymphocyte counts, and neutrophil counts. The majority of infections and blood side effects were mild to moderate in severity. Your health care provider should check you for infections before starting treatment, during treatment, and after treatment with VYVGART. Tell your health care provider if you have any history of infections. Tell your health care provider right away if you have signs or symptoms of an infection during treatment with VYVGART such as fever, chills, frequent and/or painful urination, cough, pain and blockage of nasal passages/sinus, wheezing, shortness of breath, fatigue, sore throat, excess phlegm, nasal discharge, back pain, and/or chest pain.
- **Undesirable immune reactions (hypersensitivity reactions).** VYVGART can cause the immune system to have undesirable reactions such as rashes, swelling under the skin, and shortness of breath. In clinical studies, the reactions were mild or moderate and occurred within 1 hour to 3 weeks of administration, and the reactions did not lead to VYVGART discontinuation. Your health care provider should monitor you during and after treatment and discontinue VYVGART if needed. Tell your health care provider immediately about any undesirable reactions.

Before taking VYVGART, tell your health care provider about all of your medical conditions, including if you:

- Have a history of infection or you think you have an infection
- Have received or are scheduled to receive a vaccine (immunization). Discuss with your health care provider whether you need to receive age-appropriate immunizations before initiation of a new treatment cycle with VYVGART. The use of vaccines during VYVGART treatment has not been studied, and the safety with live or live-attenuated vaccines is unknown. Administration of live or live-attenuated vaccines is not recommended during treatment with VYVGART.
- Are pregnant or plan to become pregnant and are breastfeeding or plan to breastfeed.

Tell your health care provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the common side effects of VYVGART?

The most common side effects of VYVGART are respiratory tract infection, headache, and urinary tract infection.

These are not all the possible side effects of VYVGART. Call your doctor for medical advice about side effects. You may report side effects to the US Food and Drug Administration at 1-800-FDA-1088.

What is VYVGART?

VYVGART is a prescription medicine used to treat a condition called generalized myasthenia gravis, which causes muscles to tire and weaken easily throughout the body, in adults who are positive for antibodies directed toward a protein called acetylcholine receptor (anti-AChR antibody positive).

Please see the full Prescribing Information for VYVGART and talk to your doctor.



VYVGART is a registered trademark of argenx.
For U.S. audiences only.
©2023 argenx US-EFG-21-00166 V3 06/2023

VYVGART®
(efgartigimod alfa-fcab)
Injection for Intravenous Use
400 mg/20 mL vial