

VYVGART® Hytrulo
(efgartigimod alfa and
hyaluronidase-qvfc)
Subcutaneous Injection
180 mg/mL and 2000 U/mL vial

VYVGART®
(efgartigimod alfa-fcab)
Injection for Intravenous Use
400 mg/20 mL vial

FDA-approved treatments for adults
with anti-AChR antibody positive
generalized myasthenia gravis (gMG)

Understanding your gMG symptoms: Limb weakness

AChR=acetylcholine receptor; gMG=generalized myasthenia gravis



What is generalized myasthenia gravis (gMG)?



A **rare, autoimmune, neuromuscular condition** that causes muscle weakness and fatigue

Estimated to affect
as many as
60,000
people in the U.S.

~85%
of people living with
gMG have **anti-AChR
antibody positive gMG**

Some people living with gMG
may experience limb weakness.
This may include weakness in:



Neck



Arms



Legs



Hands

By understanding and tracking
your symptoms, you can get a sense
of how they change over time.



Talk to your doctor about your symptom score on the Myasthenia Gravis Activities of Daily Living (MG-ADL) scale and share what you want to work towards with your treatment plan.

What is the MG-ADL scale? It's a tool that helps you measure the impact of gMG on daily functions. You assign a score to 8 common signs or symptoms of gMG based on how you're feeling that day. Scores range from 0 to 3—a score of 0 represents normal function and a score of 3 represents the loss of ability to perform that function. Total scores range from 0 to 24 points, with a higher score showing more severe gMG.

MG-ADL=Myasthenia Gravis Activities of Daily Living
Visit [VYVGART.com/glossary](https://www.vyvgart.com/glossary) for a glossary of terms.

This is 1 of 4 guides about different groups of gMG symptoms, and is for educational purposes only. VYVGART for intravenous (IV) infusion and VYVGART Hytrulo for subcutaneous injection have not been proven to treat a specific symptom or group of symptoms.

Find all 4 guides at
[gMGSymptoms.com](https://www.gMGSymptoms.com)

SELECT IMPORTANT SAFETY INFORMATION

Do not use VYVGART if you have a serious allergy to efgartigimod alfa or any of the other ingredients in VYVGART. Do not use VYVGART HYTRULO if you have a serious allergy to efgartigimod alfa, hyaluronidase, or any of the other ingredients in VYVGART HYTRULO. VYVGART and VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

Please see Important Safety Information throughout. Please see the full Prescribing Information for **VYVGART for IV infusion** and **VYVGART HYTRULO for subcutaneous injection**.

If you're experiencing limb weakness symptoms, you're not alone

What are some symptoms related to limb weakness?

Some people may experience difficulty with:

- Lifting their arms over their head
- Getting up from a sitting position
- Walking long distances
- Gripping heavy objects
- Climbing stairs

Please let your doctor know if you are experiencing any of these symptoms.

These symptoms
**aren't the same
for everyone**

“My kids love playing games on the floor, but if it's too hard to get down and back up again, I can't join them.”

Niki*

*Living with
anti-AChR
antibody
positive gMG*



Looking for ways to discuss your gMG symptoms and treatment goals with your doctor?



Find resources on [VYVGART.com](https://www.vyvgart.com)

SELECT IMPORTANT SAFETY INFORMATION

VYVGART and VYVGART HYTRULO may cause serious side effects, including infection.

VYVGART and VYVGART HYTRULO may increase the risk of infection. The most common infections for efgartigimod alfa-fcab-treated patients were urinary tract and respiratory tract infections. Signs or symptoms of an infection may include fever, chills, frequent and/or painful urination, cough, pain and blockage of nasal passages/sinus, wheezing, shortness of breath, fatigue, sore throat, excess phlegm, nasal discharge, back pain, and/or chest pain.

Please see Important Safety Information throughout. Please see the full Prescribing Information for **VYVGART for IV infusion** and **VYVGART HYTRULO for subcutaneous injection**.




How do your gMG symptoms affect the things that matter most to you?

Is there something you would get back to doing if your gMG symptoms and daily abilities could be improved? Talking about your goals and your gMG symptoms with your doctor may help inform your treatment plan.

“I’m very open with my doctor. I have told him I wanted to be able to go out on short hikes with my dogs again.”

Katie*

Living with anti-AChR antibody positive gMG



Download the goal-setting worksheet to help you define a Personal Ability Goal and share it with your doctor

You can also use the interactive symptom tracking tool to get your Myasthenia Gravis Activities of Daily Living (MG-ADL) symptom score.

SELECT IMPORTANT SAFETY INFORMATION

VYVGART and VYVGART HYTRULO may cause serious side effects, including allergic reactions (hypersensitivity reactions).

VYVGART and VYVGART HYTRULO can cause allergic reactions such as rashes, swelling under the skin, and shortness of breath. Hives were also observed in patients treated with VYVGART HYTRULO. Serious allergic reactions, such as trouble breathing and decrease in blood pressure leading to fainting have been reported with efgartigimod alfa-fcab.

Please see Important Safety Information throughout. Please see the full Prescribing Information for VYVGART for IV infusion and VYVGART HYTRULO for subcutaneous injection.

IMPORTANT SAFETY INFORMATION

Do not use VYVGART if you have a serious allergy to efgartigimod alfa or any of the other ingredients in VYVGART. Do not use VYVGART HYTRULO if you have a serious allergy to efgartigimod alfa, hyaluronidase, or any of the other ingredients in VYVGART HYTRULO. VYVGART and VYVGART HYTRULO can cause serious allergic reactions and a decrease in blood pressure leading to fainting.

VYVGART and VYVGART HYTRULO may cause serious side effects, including:

- **Infection.** VYVGART and VYVGART HYTRULO may increase the risk of infection. The most common infections for efgartigimod alfa-fcab-treated patients were urinary tract and respiratory tract infections. Signs or symptoms of an infection may include fever, chills, frequent and/or painful urination, cough, pain and blockage of nasal passages/sinus, wheezing, shortness of breath, fatigue, sore throat, excess phlegm, nasal discharge, back pain, and/or chest pain.
- **Allergic Reactions (hypersensitivity reactions).** VYVGART and VYVGART HYTRULO can cause allergic reactions such as rashes, swelling under the skin, and shortness of breath. Hives were also observed in patients treated with VYVGART HYTRULO. Serious allergic reactions, such as trouble breathing and decrease in blood pressure leading to fainting have been reported with efgartigimod alfa-fcab.
- **Infusion-Related Reactions.** VYVGART and VYVGART HYTRULO can cause infusion-related reactions. The most frequent symptoms and signs reported with efgartigimod alfa-fcab were high blood pressure, chills, shivering, and chest, abdominal, and back pain.

Tell your doctor if you have signs or symptoms of an infection, allergic reaction, or infusion-related reaction. These can happen while you are receiving your VYVGART or VYVGART HYTRULO treatment or afterward. Your doctor may need to pause or stop your treatment. Contact your doctor immediately if you have signs or symptoms of a serious allergic reaction.

Before taking VYVGART or VYVGART HYTRULO, tell your doctor if you:

- take any medicines, including prescription and non-prescription medicines, supplements, or herbal medicines,
- have received or are scheduled to receive a vaccine (immunization), or
- have any allergies or medical conditions, including if you are pregnant or planning to become pregnant, or are breastfeeding.

What are the common side effects of VYVGART and VYVGART HYTRULO?

The most common side effects in efgartigimod-alfa-fcab-treated patients were respiratory tract infection, headache, and urinary tract infection. Additional common side effects with VYVGART HYTRULO are injection site reactions, including rash, redness of the skin, itching sensation, bruising, pain, and hives.

These are not all the possible side effects of VYVGART and VYVGART HYTRULO. Call your doctor for medical advice about side effects. You may report side effects to the US Food and Drug Administration at 1-800-FDA-1088.

What is VYVGART® (efgartigimod alfa-fcab) for intravenous (IV) infusion and what is VYVGART® HYTRULO (efgartigimod alfa and hyaluronidase-qvfc) for subcutaneous injection?

VYVGART and VYVGART HYTRULO are both prescription medicines, each used to treat a condition called generalized myasthenia gravis, which causes muscles to tire and weaken easily throughout the body, in adults who are positive for antibodies directed toward a protein called acetylcholine receptor (anti-AChR antibody positive).

Please see the full Prescribing Information for [VYVGART](#) and the full Prescribing Information for [VYVGART HYTRULO](#), and talk to your doctor.

*Niki and Katie are paid spokespeople of argenx.

VYVGART is a registered trademark of argenx.
VYVGART Hytrulo is a trademark of argenx.
For U.S. audiences only.
©2024 US-VYV-23-00494 V1 07/2024

