

VYVGART[®]
(efgartigimod alfa-fcab)
Injection for Intravenous Use
400 mg/20 mL vial

VYVGART[®] Hytrulo
(efgartigimod alfa and
hyaluronidase-qvfc)
Subcutaneous Injection
180 mg/mL and 2000 U/mL vial

Patient Emergency Contact Card

If you need emergency medical treatment, a contact card can inform doctors and nurses that you are being treated with VYVGART/ VYVGART Hytrulo along with any other medications. This card includes vital information that may help them give you the best possible care.

Please print out this card, fill out the information, and carry it with you.

Print, cut out, fold in half, and tape shut.

! Emergency contact card !	! Emergency contact card !
My Name: _____ Medical Condition(s): _____ Emergency Contact Name: _____ Emergency Contact Number: _____ My Doctor's Name: _____ My Doctor's Number: _____	Current Medications/Supplements: (Circle one) VYVGART/VYVGART HYTRULO/ _____ _____ Allergies/Drug Allergies: _____  VYVGART is a registered trademark of argenx. For U.S. audiences only. ©2023 argenx US-VYV-23-00166 V1 09/2023

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about VYVGART® (efgartigimod alfa-fcab) for intravenous (IV) infusion and VYVGART® HYTRULO (efgartigimod alfa and hyaluronidase-qvfc) for subcutaneous injection?

VYVGART and VYVGART HYTRULO may cause serious side effects, including:

- **Infection.** VYVGART and VYVGART HYTRULO may increase the risk of infection. The most common infections for efgartigimod alfa-fcab-treated patients were urinary tract and respiratory tract infections. More patients on efgartigimod alfa-fcab vs placebo had below normal levels for white blood cell counts, lymphocyte counts, and neutrophil counts. The majority of infections and observed lower white blood cell counts were mild to moderate in severity. Your healthcare provider should check you for infections before starting treatment, during treatment, and after treatment with VYVGART or VYVGART HYTRULO. Tell your healthcare provider if you have any history of infections. Tell your healthcare provider right away if you have signs or symptoms of an infection during treatment with VYVGART or VYVGART HYTRULO such as fever, chills, frequent and/or painful urination, cough, pain and blockage of nasal passages/sinus, wheezing, shortness of breath, fatigue, sore throat, excess phlegm, nasal discharge, back pain, and/or chest pain. If a serious infection occurs, your doctor will treat your infection and may even stop your VYVGART or VYVGART HYTRULO treatment until the infection has resolved.
- **Undesirable immune reactions (hypersensitivity reactions).** VYVGART and VYVGART HYTRULO can cause the immune system to have undesirable reactions such as rashes, swelling under the skin, and shortness of breath. Hives were also observed in patients treated with VYVGART HYTRULO. In clinical studies, the reactions were mild or moderate and occurred within 1 hour to 3 weeks of administration, and the reactions did not lead to treatment discontinuation. Your healthcare provider should monitor you during and after treatment and discontinue VYVGART or VYVGART HYTRULO if needed. Tell your healthcare provider immediately about any undesirable reactions to VYVGART or VYVGART HYTRULO.

Before taking VYVGART or VYVGART HYTRULO, tell your healthcare provider about all of your medical conditions, including if you:

- Have a history of infection or you think you have an infection.
- Have received or are scheduled to receive a vaccine (immunization). Discuss with your healthcare provider whether you need to receive age-appropriate immunizations before initiation of a new treatment cycle with VYVGART or VYVGART HYTRULO. The use of vaccines during treatment with VYVGART or VYVGART HYTRULO has not been studied, and the safety with live or live-attenuated vaccines is unknown. Administration of live or live-attenuated vaccines is not recommended during treatment with VYVGART or VYVGART HYTRULO.
- Are pregnant or plan to become pregnant and are breastfeeding or plan to breastfeed.

Tell your health care provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the common side effects of VYVGART and VYVGART HYTRULO?

The most common side effects of efgartigimod-alfa-fcab-treated patients were respiratory tract infection, headache, and urinary tract infection. Additional common side effects with VYVGART HYTRULO are injection site reactions, including rash, redness of the skin, itching sensation, bruising, pain, and hives.

These are not all the possible side effects of VYVGART and VYVGART HYTRULO. Call your doctor for medical advice about side effects. You may report side effects to the US Food and Drug Administration at 1-800-FDA-1088.

What is VYVGART for IV infusion and what is VYVGART HYTRULO for subcutaneous injection?

VYVGART and VYVGART HYTRULO are both prescription medicines, each used to treat a condition called generalized myasthenia gravis, which causes muscles to tire and weaken easily throughout the body, in adults who are positive for antibodies directed toward a protein called acetylcholine receptor (anti-AChR antibody positive).

Please see the full Prescribing Information for [VYVGART](#) and the full Prescribing Information for [VYVGART HYTRULO](#) and talk to your doctor.